

[Web](#) [Images](#) [Videos](#) [Maps](#) [News](#) [Shopping](#) [Gmail](#) [more ▼](#)

[Sign in](#)

Google scholar

((foldback or folded or "fold over" or hairpin) an

Search

[Advanced Scholar Search](#)  
[Scholar Preferences](#)

Scholar

Articles and patents

anytime

include citations



Create email alert

Results 1 - 10 of about 23. (0.17 sec)

[CITATION] Organization and replication of parvovirus DNA

Ki Bems, WW Hauswirth - Organization and replication of viral DNA, 1982

[Cited by 12](#) - [Related articles](#)

[Rescue of the adeno-associated virus genome from a plasmid vector: evidence for rescue by replication](#)

[nih.gov \[HTML\]](#)

P Ward, P Elias, RM Linden - Journal of virology, 2003 - Am Soc Microbiol

... The **hairpin** primers can be recreated indefinitely by the process designated terminal resolution, which consists of ... The duplex ITR can then **fold over** and serve as a **primer** for full-length ... copy of the AAV genome is replicated from the plasmid by a **fold-back** replication mechanism ...

[Cited by 5](#) - [Related articles](#) - [BL Direct](#) - [All 10 versions](#)

[De novo synthesis of RNA by the dengue virus RNA-dependent RNA polymerase exhibits temperature dependence at the initiation but not elongation phase](#)

[jbc.org \[HTML\]](#)

M Ackermann, R Padmanabhan - Journal of Biological Chemistry, 2001 - ASEMB

... either positive (5'-AGCTGTACGATGGCGTAG-3') or negative strand (5'-CTACGCCATCGTACAGCT-3') **primer** and were ... agarose gel system: a template-sized RNA (1×) and a **hairpin** RNA, twice ... dimer species could also have resulted by 3'-elongation of a "**folded-back**" structure ...

[Cited by 123](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

[G-quadruplex DNA: a potential target for anti-cancer drug design](#)

[cell.com \[HTML\]](#)

H Han, LH Hurley - Trends in pharmacological sciences, 2000 - Elsevier

... One model proposes that the single-stranded G-rich overhang might **fold back** to form a ... During the translocation, a **fold-over** G-quadruplex is transiently formed from telomeric DNA to provide the ... newly synthesized G-rich strand forms a secondary structure such as a **hairpin** or G ...

[Cited by 157](#) - [Related articles](#) - [All 2 versions](#)

[Role of miRNA in carcinogenesis and biomarker selection: a methodological view](#)

FE Ahmed - Expert Review of Molecular Diagnostics, 2007 - ingentaconnect.com

... Drosha and a dsDNA binding protein DGCR8/Pasha) recognize the distinct **fold-back** (**hairpin**) secondary structure ... as a similarity measure for 29 global and intrin- sic **hairpin** folding attributes ... Trained on 200 human pre- miRNAs and 400 pseudo **hairpins**, miPred achieved 93.50 ...

[Cited by 14](#) - [Related articles](#) - [BL Direct](#) - [All 5 versions](#)

[HTML] [Ago2 immunoprecipitation identifies predicted MicroRNAs in human embryonic stem cells and neural precursors](#)

[nih.gov \[HTML\]](#)

LA Goff, J Davila, MR Swerdel, JC Moore, RI Cohen, H ... - 2009 - ncbi.nlm.nih.gov

... Master Mix, Applied Biosystems) using one microRNA-specific **primer** and one **primer** specific for the ... These precursors were **folded** with RNAFold [40] to identify characteristic **hairpin** structures. ... and those predicted microRNAs that were enriched at least 4-**fold over** IgG controls ...

[Cited by 9](#) - [Related articles](#) - [All 7 versions](#)

[Recombinant viral vector system](#)

RJ Samulski, X Xiao - US Patent 5,478,745, 1995 - Google Patents

... regions are two 145 nucleotide inverted terminal (ITR) repeat sequences that contain palindromic sequences that can **fold over** to 60 form **hairpin** structures that ... which are unique in that they can not only basepair with each other, but also individually **fold back** on themselves ...

[Cited by 2](#) - [Related articles](#) - [All 5 versions](#)

[Targeting human telomerase in cancer therapy](#)

S Huard, C Autexier - Current Medicinal Chemistry-Anti- ... - 2002 - ingentaconnect.com

... of telomeres can form a **hairpin** structure and dimerize with a similar **hairpin** from a ... Another model suggests that the 3'-overhang can **fold over** to form an intramolecular G- quadruplex ... have shown that the single-stranded 3'-overhang at telomeres can **fold back** and invade the ...

[Cited by 13](#) - [Related articles](#) - [BL Direct](#) - [All 2 versions](#)

[PDF] [Ago2 Immunoprecipitation Identifies Predicted MicroRNAs in Human Embryonic Stem](#)

[systembio.com \[PDF\]](#)

LA Goff, J Davila, MR Swerdel, JC Moore, RI Cohen ... - 2009 - systembio.com

... Green Master Mix, Applied Biosystems) using one microRNA-specific **primer** and one **primer** specific for ... the wiggle tracks) and the solid line represents the remaining predicted **hairpin** precursor. ... and those predicted microRNAs that were enriched at least 4-**fold over** IgG controls ...

[Related articles](#) - [View as HTML](#) - [All 4 versions](#)

[PDF] [REKOMBINANTES VIRALES VEKTORSYSTEM SYSTEME PORTEUR VIRAL DE RECOMBINAISON](#)

[epo.org \[PDF\]](#)

A ES, F LI, LUMC NL, PT SE - GENE, 1992 - data.epo.org

... regions are two 145 nucleotide inverted terminal (ITR) repeat sequences that contain palindromic sequences that can **fold over** to form **hairpin** structures that ... which are unique in that they can not only basepair with each other, but also individually **fold back** on themselves ...

[View as HTML](#)